

## Assessment of Tumor-infiltrating Lymphocytes Predicts the Behavior of Early-stage Oral Tongue Cancer

Heikkinen, Ilkka

2019-10

---

Heikkinen , I , Bello , I O , Wahab , A , Hagström , J , Haglund , C , Coletta , R D , Nieminen , P , Makitie , A A , Salo , T , Leivo , I & Almangush , A 2019 , ' Assessment of Tumor-infiltrating Lymphocytes Predicts the Behavior of Early-stage Oral Tongue Cancer ' , American Journal of Surgical Pathology , vol. 43 , no. 10 , pp. 1392-1396 . <https://doi.org/10.1097/PAS.0000000000000000>

---

<http://hdl.handle.net/10138/324427>

<https://doi.org/10.1097/PAS.00000000000001323>

---

cc\_by\_nc

acceptedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

**Assessment of tumor-infiltrating lymphocytes predicts the behavior of early-stage oral tongue cancer**

**Ilkka Heikkinen, DDS,<sup>a</sup> Ibrahim O. Bello, DDS, PhD,<sup>b</sup> Awais Wahab, DDS,<sup>c</sup> Jaana Hagström, DDS, PhD,<sup>d</sup> Caj Haglund, MD, PhD,<sup>e</sup> Ricardo D. Coletta, DDS, PhD,<sup>f</sup> Pentti Nieminen, MSc, PhD,<sup>g</sup> Antti A. Mäkitie, MD, PhD,<sup>h</sup> Tuula Salo, DDS, PhD,<sup>i\*</sup> Ilmo Leivo, MD, PhD,<sup>j\*</sup> Alhadi Almangush, DDS, PhD<sup>k\*</sup>**

<sup>a</sup>Department of Pathology, University of Helsinki, Helsinki, Finland.

Department of Oral and Maxillofacial Diseases, University of Helsinki, Helsinki, Finland.

<sup>b</sup>Department of Oral Medicine and Diagnostic Sciences, King Saud University College of Dentistry, Riyadh, Saudi Arabia.

Department of Pathology, University of Helsinki, Helsinki, Finland.

<sup>c</sup>Department of Pathology, University of Helsinki, Helsinki, Finland.

Department of Oral and Maxillofacial Diseases, University of Helsinki, Helsinki, Finland.

<sup>d</sup>Department of Pathology, University of Helsinki, Helsinki, Finland.

Research Programs Unit, Translational Cancer Biology, University of Helsinki, Helsinki, Finland.

<sup>e</sup> Research Programs Unit, Translational Cancer Biology, University of Helsinki, Helsinki, Finland.

Department of Surgery, University of Helsinki and Helsinki University Hospital, Helsinki, Finland.

<sup>f</sup> Department of Oral Diagnosis, School of Dentistry, University of Campinas, Piracicaba, São Paulo, Brazil.

<sup>g</sup> Medical Informatics and Data Analysis Research Group, University of Oulu, Oulu, Finland.

<sup>h</sup> Department of Otorhinolaryngology – Head and Neck Surgery, University of Helsinki and Helsinki University Hospital, Helsinki, Finland.

Research Programme in Systems Oncology, Faculty of Medicine, University of Helsinki, Helsinki, Finland.

Division of Ear, Nose and Throat Diseases, Department of Clinical Sciences, Intervention and Technology,

1   **Funding:** This study funded by Turku University Hospital Fund, Helsinki University Hospital  
2   Research Fund, the Finnish Dental Society, the Rauha Ahokas Foundation, K. Albin Johanssons  
3   Foundation, the Sigrid Juselius Foundation, the Finnish Cancer Society, Finska Läkaresällskapet, and  
4   the Maritza and Reino Salonen Foundation.

5

6   **Conflict of interest:** None declared.

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

1   **Abstract**

2   Tumor-infiltrating lymphocytes (TILs) have shown a promising prognostic value in many epithelial  
3   cancers. We sought to assess the prognostic value of TILs in a multicenter cohort of early oral tongue  
4   squamous cell carcinoma (OTSCC). The percentage of TILs was assessed on the surgical resection  
5   slides stained with hematoxylin and eosin (HE). The assessment of TILs was performed in the stromal  
6   compartment and in the intra-epithelial compartment (at the invasive front and at the center of the  
7   tumor). We followed the method that was described recently by the International Immuno-Oncology  
8   Biomarker Working Group for the assessment of TILs. A total of 308 cases from the five Finnish  
9   university hospitals and from A.C. Camargo Cancer Center, São Paulo, Brazil were included. We  
10   found a promising prognostic value for stromal TILs at the invasive front in the multivariable analysis  
11   with a hazard ratio of 2.61 (95%CI 1.77-3.83;  $P<0.001$ ) for overall survival, 1.99 (95%CI 1.07-3.69;  
12    $P=0.040$ ) for disease-specific survival, and 1.94 (95%CI 1.14-3.29;  $P=0.020$ ) for disease-free  
13   survival. In conclusion, evaluation of TILs is simple and can aid in identifying the high-risk cases of  
14   early OTSCC. The method introduced by the International Immuno-Oncology Biomarker Working  
15   Group can be used for standardized determination of TILs in early OTSCC.

16  
17   **Keywords:** Oral tongue squamous cell carcinoma (OTSCC); Tumor-infiltrating lymphocytes (TILs);  
18   Overall survival (OS); Disease-specific survival (DSS); Disease-free survival (DFS).

19  
20  
21  
22  
23  
24  
25  
26

## 1    **Introduction**

2    Oral tongue squamous cell carcinoma (OTSCC) is associated with worse survival compared to SCC  
3    of other sites of the oral cavity <sup>1</sup>. The current prognostic biomarkers of early-stage OTSCC do not  
4    provide optimal risk stratification <sup>2</sup>. Therefore, searching for "informative, simple and reliable"  
5    biomarkers is of great importance, since those biomarkers will form a cornerstone for individualized  
6    treatment. The role of tumor microenvironment in cancer development and prognosis has evoked  
7    increasing interest among cancer researchers in recent years. Immune cells comprise one component  
8    of the tumor microenvironment and the immune infiltrates associating with tumors have been studied  
9    widely <sup>3,4</sup>.

Pathological prognostication of clinical behavior is usually based on

tools or antibodies, and easy incorporation into standard pathology report. In this study, we investigated the significance of TILs as a potential prognostic marker in early OTSCC using a multi-institutional cohort that we have used in our previous study <sup>9</sup>.

## Material and Methods

**Patients:** This study included 308 patients (163 men and 145 women) treated for early OTSCC at one of the five Finnish University Hospitals (Helsinki, Turku, Tampere, Oulu, and Kuopio) or at the A.C. Camargo Cancer Center, São Paulo, Brazil. The primary treatment of patients in our cohort was surgical resection. The use of patient samples and the data enquiry were approved by the above-mentioned University hospitals, by the Brazilian Human Research Ethics Committee and by the Finnish National Supervisory Authority for Welfare and Health (VALVIRA).

**Scoring of TILs:** A training session was arranged to familiarize the observers with the scoring criteria. We evaluated TILs according to a scoring method introduced recently by the International Immuno-Oncology Biomarkers Working Group <sup>7</sup>. In brief, intra-tumoral TILs were scored as the percentage of tumor islands occupied by lymphocytes; and stromal TILs were defined as the percentage of stroma occupied by lymphocytes. Any stromal area that did not relate directly to the tumor was not included in the estimation of TILs. Moreover, areas of fibrosis or central necrosis were not included in the assessment of TILs <sup>7</sup>. The percentage of TILs was assessed in two regions of each sample (at the invasive front and at the central part of the tumor). Assessment of TILs was made in two compartments, the intra-tumoral (i.e. intra-epithelial) and the stromal compartment. Tumor slides were visually scanned by light microscope, and the percentage of TILs at the invasive front was estimated separately for the intra-tumoral part and for the stromal part. The same scoring method was used to assess TILs in the central region of the tumor. The TILs working group guidelines <sup>7, 10</sup> has pointed out “*Do not focus on hot spots*”. Therefore, the average of TILs in the stromal area was considered when reporting stromal TILs. Similarly, the average of TILs in the tumor area was

1 considered when reporting intra-tumoral TILs. The percentage of TILs was assessed semi-  
2 quantitatively as an incremental parameter (e.g. 5%, 10%, 20%, 30%...) as previously described <sup>7, 10</sup>.  
3 For example, 50% stromal TILs indicates that half of the stromal area is occupied by infiltrating  
4 lymphocytes. The sample was scanned at low magnification (with 5-10× objectives), then the average  
5 percentage of TILs across microscopic fields was estimated at higher magnification (with 20-40×  
objectives). At least five fields were evaluated to assess the average of TILs.





Intra-tumoral infiltrating lymphocytes did not show a significant prognostic value in univariable analyses of OS (HR 1.14, 95%CI 0.76-1.68;  $P = 0.51$ ), DSS (HR 1.06, 95%CI 0.57-1.94;  $P = 0.863$ ) or DFS (HR 1.06, 95%CI 0.63-1.78;  $P = 0.834$ ). The insignificant prognostic values of intra-tumoral infiltrating lymphocytes were confirmed in the multivariable analyses of OS (HR 1.05, 0.71-1.55;  $P = 0.818$ ), DSS (HR 1.19, 95%CI 0.63-2.22;  $P = 0.594$ ) and DFS (HR 1.09, 95%CI 0.64-1.85;  $P = 0.761$ ).

## Discussion

For clinicians, it is a dilemma to identify those cases of early OTSCC that will potentially have an aggressive behavior and would thus benefit from chemoradiotherapy and/or neck dissection. Many factors will determine tumor behavior and risk of poor survival. Of these factors, immune response has been shown to possess a fundamental role in identifying aggressive tumors<sup>6</sup>. In this study, we used the HE-stained slides from surgical specimens to evaluate TILs in a multicenter series of early OTSCC and we found that tumors with low percentage of stromal TILs ( $\leq 20\%$ ) at the invasive front have a significantly poor survival (overall, disease-specific and disease-free).

As the current method of evaluating TILs was based on the use of routine H-E slides, it will be possible to assess them routinely in the daily practice of pathologists and will provide a better histopathologic prognostication based on immune response. Such prognostication may be useful for clinical decision-making including, for example, selecting early-stage OTSCC cases for multimodality treatments. Recent research highlighted the role of the immune cells in modulating cancer invasion and metastasis<sup>12</sup>. Therefore, the immune response is assumed to influence the clinical behavior of tumors. In fact, tumors of the same clinical stage and/or same histopathologic grade may have extremely variable immune responses<sup>6</sup>. Thus, the immunological heterogeneity of early OTSCC can be utilized to classify cases into low-risk and high-risk groups.

1        Recently, we have systematically reviewed the prognostic value of all immune checkpoints  
2        that have been studied in oral squamous cell carcinoma <sup>13</sup>. We noted that the currently available body  
3        of evidence still require further research and none of the studied immune biomarkers can be approved  
4        for prognostication in daily practice <sup>13</sup>. In addition, previous studies have used a specific  
5        immunohistochemical staining which is not routinely ordered by the pathologists. In our previous  
6        research, we evaluated the lymphocytic host response (LHR) according to criteria described in the  
7        histologic risk score <sup>14</sup> and we reported a low prognostic value of that criteria for prognostication of  
8        early OTSCC <sup>15</sup>. Noteworthy, LHR was divided into three categories (strong, intermediate or weak)  
9        without quantitative measurement of the immune response. On the other hand, the International  
10        Immuno-Oncology Biomarkers Working Group has indicated that assessment of TILs should be done  
11        as a continuous semiquantitative variable and did not determine risk threshold between high and low  
12        TILs in early OTSCC <sup>7</sup>.

13        Currently, a method for overall assessment of TILs (i.e. using HE-stained slides) after  
14        neoadjuvant treatment of OTSCC has not yet been established. In cases with neoadjuvant therapy  
15        determining the area of the residual tumor to be used for evaluation of TILs requires further research  
16        <sup>16</sup>. A small and/or superficial preoperative biopsy of oral cancer might also be a problematic as it may  
17        not include the most important area for assessment of TILs (i.e. within the borders of the invasive  
18        tumor). Such limitation of biopsies of oral cancer has been noted during the assessment of other  
19        prognostic features <sup>17</sup>. Therefore, it is necessary to consider having a representative sample that is  
20        deep enough including the invasive front for the assessment of TILs.

21        Recently, there is a worldwide research effort to standardize determination of TILs. Many  
22        studies have reported that a strong lymphocytic infiltration associates with favorable clinical outcome  
23        of different tumors including those of head and neck <sup>18-20</sup>. Such favorable outcome could be due to  
24        the destruction of cancer cells and anti-tumor effect of the immune response <sup>21</sup>. In conclusion,  
25        assessment of TILs has a reliable prognostic value in early OTSCC. The method introduced recently

1 by the International Immuno-Oncology Biomarkers Working Group is simple, cost-effective and can  
 2 be easily included in the pathology report. To the best of our knowledge, this is the first multicenter  
 3 study on early OTSCC applying the new criteria for evaluation of TILs. Our finding on TILs can be  
 4 a key step towards the routine measurement of immune-response in early OTSCC and that could  
 5 enhance the personalized treatment approach by classifying risk groups based on immune response.

6  
 7

## 8 **References**

- 9 1. Farhood Z, Simpson M, Ward GM, et al. Does anatomic subsite influence oral cavity cancer  
 10 mortality? A SEER database analysis. *Laryngoscope*. 2018.
- 11 2. Almangush A, Heikkinen I, Makitie AA, et al. Prognostic biomarkers for oral tongue  
 12 squamous cell carcinoma: a systematic review and meta-analysis. *Br J Cancer*. 2017;117:856-866.
- 13 3. Wang M, Zhao J, Zhang L, et al. Role of tumor microenvironment in tumorigenesis. *J Cancer*.  
 14 2017;8:761-773.
- 15 4. Giraldo NA, Sanchez-Salas R, Peske JD, et al. The clinical role of the TME in solid cancer.  
 16 *Br J Cancer*. 2019;120:45-53.
- 17 5. Pandya PH, Murray ME, Pollok KE, et al. The Immune System in Cancer Pathogenesis:  
 18 Potential Therapeutic Approaches. *J Immunol Res*. 2016;2016:4273943.
- 19 6. Galon J, Pages F, Marincola FM, et al. Cancer classification using the Immunoscore: a  
 20 worldwide task force. *J Transl Med*. 2012;10:205.
- 21 7. Hendry S, Salgado R, Gevaert T, et al. Assessing Tumor-Infiltrating Lymphocytes in Solid  
 22 Tumors: A Practical Review for Pathologists and Proposal for a Standardized Method from the  
 23 International Immuno-Oncology Biomarkers Working Group: Part 2: TILs in Melanoma,  
 24 Gastrointestinal Tract Carcinomas, Non-Small Cell Lung Carcinoma and Mesothelioma, Endometrial  
 25 and Ovarian Carcinomas, Squamous Cell Carcinoma of the Head and Neck, Genitourinary  
 26 Carcinomas, and Primary Brain Tumors. *Adv Anat Pathol*. 2017;24:311-335.
- 27 8. Dieci MV, Radosevic-Robin N, Fineberg S, et al. Update on tumor-infiltrating lymphocytes  
 28 (TILs) in breast cancer, including recommendations to assess TILs in residual disease after  
 29 neoadjuvant therapy and in carcinoma in situ: A report of the International Immuno-Oncology  
 30 Biomarker Working Group on Breast Cancer. *Semin Cancer Biol*. 2018;52:16-25.
- 31 9. Almangush A, Heikkinen I, Bakhti N, et al. Prognostic impact of tumour-stroma ratio in early-  
 32 stage oral tongue cancers. *Histopathology*. 2018;72:1128-1135.
- 33 10. Salgado R, Denkert C, Demaria S, et al. The evaluation of tumor-infiltrating lymphocytes  
 34 (TILs) in breast cancer: recommendations by an International TILs Working Group 2014. *Ann Oncol*.  
 35 2015;26:259-271.
- 36 11. Kleinbaum DG, Klein M. Survival Analysis: A Self-Learning Text, Third Edition. 2012.
- 37 12. Kitamura T, Qian BZ, Pollard JW. Immune cell promotion of metastasis. *Nat Rev Immunol*.  
 38 2015;15:73-86.
- 39 13. Sievilainen M, Almahmoudi R, Al-Samadi A, et al. The Prognostic Value of Immune  
 40 Checkpoints in Oral Squamous Cell Carcinoma. *Oral Dis*. 2018.

- 1 14. Brandwein-Gensler M, Smith RV, Wang B, et al. Validation of the histologic risk model in a  
2 new cohort of patients with head and neck squamous cell carcinoma. *Am J Surg Pathol*. 2010;34:676-  
3 688.
- 4 15. Almangush A, Bello IO, Keski-Santti H, et al. Depth of invasion, tumor budding, and worst  
5 pattern of invasion: prognostic indicators in early-stage oral tongue cancer. *Head Neck*. 2014;36:811-  
6 818.
- 7 16. Kurozumi S, Inoue K, Matsumoto H, et al. Prognostic utility of tumor-infiltrating  
8 lymphocytes in residual tumor after neoadjuvant chemotherapy with trastuzumab for HER2-positive  
9 breast cancer. *Sci Rep*. 2019;9:1583.
- 10 17. Dhanda J, Uppal N, Chowlia H, et al. Features and prognostic utility of biopsy in oral  
11 squamous cell carcinoma. *Head Neck*. 2016;38 Suppl 1:E1857-1862.
- 12 18. Denkert C, von Minckwitz G, Darb-Esfahani S, et al. Tumour-infiltrating lymphocytes and  
13 prognosis in different subtypes of breast cancer: a pooled analysis of 3771 patients treated with  
14 neoadjuvant therapy. *Lancet Oncol*. 2018;19:40-50.
- 15 19. Rakaee M, Kilvaer TK, Dalen SM, et al. Evaluation of tumor-infiltrating lymphocytes using  
16 routine H&E slides predicts patient survival in resected non-small cell lung cancer. *Hum Pathol*.  
17 2018;79:188-198.
- 18 20. Almangush A, Ruuskanen M, Hagstrom J, et al. Tumor-infiltrating lymphocytes associate  
19 with outcome in non-endemic nasopharyngeal carcinoma: a multicenter study. *Hum Pathol*. 2018.
- 20 21. Fridman WH, Pages F, Sautes-Fridman C, et al. The immune contexture in human tumours:  
21 impact on clinical outcome. *Nat Rev Cancer*. 2012;12:298-306.
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34

1 **Figure legend**

2 **Figure 1:** Tumor-infiltrating lymphocytes (TILs) were evaluated on full sections of early OTSCC  
3 stained with hematoxylin and Eosin. Representative samples classified as low ( $\leq 20\%$ ) stromal TILs  
4 (A); and high ( $> 20\%$ ) stromal TILs (B).

5  
6 **Figure 2:** Kaplan-Meier curves for stromal TILs with OS (A), DSS (B), and DFS (C).

7 TILs: Tumor-infiltrating lymphocytes; OS: Overall survival; DSS: Disease-specific survival; DFS:  
8 Disease-free survival.

9









